



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.


Let $OB=OC=OD=s$, $OI=x$, $OG=OH=w$; then $OA=s\sqrt{2}$, area of $\square ABOC=s^2$, and area of circle radius of which is $OA=2\pi s^2$.

First Solution.—Knowing the favorable chances and the whole number of chances, the required chance becomes

$$C=\frac{s^2}{2\pi s^2}=\frac{1}{2\pi}\dots\dots(A).$$

SECOND SOLUTION.

The arc JIK increases for all values of OI less than OD ; and the arc HG decreases for all values of OG greater than OD . The required chance, therefore, becomes



$$\begin{aligned} C &= \frac{1}{2\pi s^2} \left\{ \frac{\pi}{2} \int_0^s x dx + \int_s^{s\sqrt{2}} \left[\frac{\pi}{2} - 2 \cos^{-1} \left(\frac{s}{w} \right) \right] w dw \right\} \\ &= \frac{1}{8} + \frac{1}{2\pi^2} \left[\frac{\pi s^2}{4} - 2 \int_s^{s\sqrt{2}} \cos^{-1} \left(\frac{s}{w} \right) w dw \right] \\ &= \frac{1}{4} - \frac{1}{2\pi s^2} \left[w^2 \cos^{-1} \left(\frac{s}{w} \right) - s \sqrt{(w^2 - s^2)} \right]_s^{s\sqrt{2}} \\ &= \frac{1}{4} - \frac{1}{2\pi} \left[\frac{\pi}{2} - 1 \right] = \frac{1}{2\pi} \dots\dots(B). \end{aligned}$$

This problem was solved with different results by H. W. DRAUGHON, P. H. PHILBRICK, and WILLIAM, B. MILWARD. If we have space, some of these solutions will be published next month.

PROBLEMS.

6. Proposed by Professor J. F. W. SCHEFFER, Hagerstown, Maryland.

Find the average length of all the diameters that can be drawn in a given ellipse.

7. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

A letter received is known to have come from Oshkosh or Ashland. The only two consecutive letters legible on the postmark are SH. What is the probability that the letter received came from Oshkosh?

8. Proposed by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Prove that the mean area of all triangles having their vertices upon the surface of a given triangle and bases parallel to the base of the given triangle, is $\frac{13}{210}$ (area of given triangle).

Solutions to these problems should be received on or before July 1st.